



The Perspective of Social Determinants of Health (SDOH) for Diabetes Mellitus

Bando H^{1,2,3,*}

¹Tokushima University / Medical Research, Tokushima, Japan

²Japan Low Carbohydrate Diet Promotion Association (JLCDPA), Kyoto, Japan

³Shikoku Island Division, Integrative Medicine Japan (IMJ), Tokushima, Japan

*Corresponding author: Bando H, Tokushima University /Medical Research, Nakashowa 1-61, Tokushima 770-0943 Japan; Tel: +81-90-3187-2485; E-mail: pianomed@bronze.ocn.ne.jp

Received date: 19 April 2021; Accepted date: 24 April 2021; Published date: 29 April 2021

Citation: Bando H (2021). The Perspective of Social Determinants of Health (SDOH) for Diabetes Mellitus. SunText Rev Endocrine Care 1(1): 102.

Copyright: © 2021 Bando H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Clinical practice for diabetes mellitus requires bio-psycho-social aspects. For continuing adequate health care, the perspective of social determinants of health (SDOH) has been recognized as indispensable intervention target for evaluating health equity. SDOH include socio-economic situation, neighbourhood and physical circumstance, food environment, health care, social context with social support capital and relationship. For contextual factors, American Diabetes Association (ADA) established the writing committee and reviewed the detail of SDOH. They included Healthy People 2020, Social Determinants of Health of WHO, the County Health Rankings Model and others. Thus, enough understanding of environmental and social factors would be necessary for diabetes.

Keywords: Diabetes mellitus (DM); Social determinants of health (SDOH); American Diabetes Association (ADA); World Health Organization (WHO); LCD promotion association (JLCDPA)

Commentary Article

In recent years, non-communicable diseases (NCDs) have become widespread along with changes in medical and social structures. They include diabetes, obesity, hypertension, dyslipidemia, cerebrovascular accidents (CVA), and cardiovascular diseases (CVD). These diseases are very common and may cause various clinical complications. Among them, diabetes is especially crucial disease in the clinical setting of primary care [1].

Authors and collaborators have continued clinical practice and research for long. The overall area includes diabetes, primary care and integrative medicine. For effective regimen for NCDs, low carbohydrate diet (LCD) has been introduced to medical and health area [2]. Three types of LCD were proposed as super-LCD, standard-LCD, petite-LCD including carbohydrate ratio as 12%, 26%, 40%, respectively [3]. Furthermore, we established Japan LCD promotion association (JLCDPA) and developed LCD in Japan as medical and social movement.

During our practice and activities, an important and common clinical concept has been present. It is a bio-psycho-social perspective, and the basis for medical care in any area [4]. In this paper, social perspective on diabetes will be described.

Health care has been gradually emphasized on value-based care and population-based health outcomes. Consequently, the perspective of social determinants of health (SDOH) has been recognized as indispensable intervention target for evaluating health equity [5,6]. Most recently, crucial problem of pandemic infection has been persisting worldwide concerning the COVID-19. It has brought highlighted unequal situation related to racial minority groups and disproportional communities. In responses to these simultaneous occurrence, some proposals of SDOH have been published by professional medical organizations, including American College of Physicians, Society of General Internal Medicine, American Academy of Pediatrics, National Academy of Medicine, and other medical associations [7,8]. Further, their calls to action would focus on the improvement of the determinants at the levels of the organization, individual and policy [9].

For clinical practice for diabetes, several fundamental problems have been present such as economic costs, incidence and prevalence and disproportionate burden in the population [10]. American Diabetes Association (ADA) proposed a statement formerly on socioecological determinants of type 2 diabetes

Citation: Bando H (2021). The Perspective of Social Determinants of Health (SDOH) for Diabetes Mellitus. SunText Rev Endocrine Care 1(1): 102.

mellitus (T2DM) and prediabetes [11]. From the viewpoint of SDOH, to understand and to advance the health improvement opportunities will be obtained for diabetic patients. Consequently, ADA has convened the writing committee for diabetes toward the purpose of informing SDOH broadly. It includes the relationship with SDOH and diabetes risk/outcomes, and various literature review concerning SDOH [11].

When searching for medical literature for the keyword of SDOH, the results include several categories concerning SDOH [12]. They are i) socio-economic situation (occupation, income, education), ii) neighborhood and physical circumstance (building environment, residence condition, toxic exposure in the environment), iii) food environment (accessibility for food, insecurity for food), iv) health care region (affordability, accessibility, quality), v) social context (social support, social capital and budget, social relationship). These factors and their linkages are investigated in the light of health care and community-based perspective.

To recognize the definitions of some key words would be important. Diabetic risks and outcomes are not same but different, in which multiple contributors exist such as clinical, biological, social and other related factors [13]. Lots of scientific literatures showed the adverse influence of health disparities related to diabetes [14]. In order to investigate the matter of healthy equity, enough understanding of environmental and social factors would be required, in which 50-60 % of outcomes of health problems can be included. Those environmental and social factors have been collectively recognized as SDOH.

For contextual factors, the writing committee of ADA has reviewed the detail of SDOH in the light of terminology and classifications. They included Healthy People 2020 [15], Social Determinants of Health of WHO, the County Health Rankings Model and others. Common elements in the frameworks are economic and socioeconomic determinants as foremost. In the case of food SDOH factors, they are divided into several groups, such as material circumstance, neighborhood environment, built environment, economic stability and independent category [15].

As related to the perspective of SDOH, a statistic marker of the adherence has been observed concerning the problems of actual satisfactory intake of medicine. The number of the patients with diabetes mellitus is increasing across the world. From the data of World Health Organization (WHO) (2020), adults aged 18 yo and older had diabetes mellitus as 8.5% in 2014. An increase of 5% was observed in premature mortality from diabetes during 2000-2016. In the countries with lower-middle-income level, the premature mortality rate due to diabetes gradually increased during this period. Furthermore, diabetes became the direct cause in 2016 as 1.6million deaths and hypertension was the cause in 2012 as 2.2 million deaths worldwide [16]. High blood glucose or hyperglycemia has been the common situation of uncontrolled

diabetes, and it will bring serious complications to various systems in the body.

United States Center for Disease Control and Prevention showed the increasing number of adults diabetic patients, who take pills, insulin or both [17]. On the other hand, higher ratio of them seemed to have inadequate adherence and acceptance due to the shortage of the knowledge [18]. Consequently, sufficient information about diabetes has to be learned in order to continue optimal glucose variability with satisfactory medication adherence.

Concerning the medication adherence of diabetic patient, a review on knowledge of diabetes and actual practice was investigated [19]. For the analysis, 18 articles were included, with the result that the knowledge level did not guarantee the medication adherence for health-seeking practice. Some factors were observed that predicting the information level and actual practice are probably from sociodemographic characteristics [19].

In summary, this article introduced some clinical matters of SDOH and its related perspective concerning diabetes mellitus. Diabetes and endocrine diseases have characteristics of long period treatment, in which patient and medical staff can understand bio-psycho-social situation and cooperate together. This article will hopefully become a useful reference in the clinical diabetic practice.

Conflicts of Interest

The author declares that they have no conflicts of interest.

Funding

The author does not have any funding concerning our medical research.

References

1. American Diabetes Association. Diabetes technology: Standards of Medical Care in Diabetes. 2021. *Diabetes Care*. 2021; 44: S85-S99.
2. Yamashita H, Kato Y, Bando H, Kanazawa S, Tanaka M, Sueki E, et al. Relationship of glucose variability and daily lifestyle by continuous glucose monitoring (CGM). *Asp Biomed Clin Case Rep*. 2020; 3: 206-212.
3. Bando H. Useful tips for actual low carbohydrate diet (lcd) with super-, standard- and petit-lcd methods. *EC Nutrition* 2020; 15: 1-4.
4. Hayashi K, Yasuoka T, Bando H, Miki K, Nakagawa M, Zushi T, et al. Useful Xultophy for Older Diabetic with Various Problems. *SunText Rev Med Clin Res* 2021; 2: 126.
5. Ogunwole SM, Golden SH. Social determinants of health and structural inequities-root causes of diabetes disparities. *Diabetes Care*. 2021; 44: 11-13.



6. Chin MH. Creating the business case for achieving health equity. *J Gen Intern Med.* 2016; 31: 792-796.
7. National Academies of Sciences, Engineering, and Medicine. *Integrating Social Care into the Delivery of Health Care: Moving Upstream to Improve the Nation's Health.* Washington, DC, National Academies Press, 2019.
8. Daniel H, Bornstein SS, Kane GC. Health and Public Policy Committee of the American College of Physicians. Addressing social determinants to improve patient care and promote health equity: an American College of Physicians position paper. *Ann Intern Med.* 2018; 168: 577-578.
9. Chin MH, King PT, Jones RG. Lessons for achieving health equity comparing Aotearoa/ New Zealand and the United States. *Health Policy.* 2018; 122: 837-853.
10. Haire-Joshu D, Hill-Briggs F. The next generation of diabetes translation: a path to health equity. *Annu Rev Public Health.* 2019; 40: 391-410.
11. Hill JO, Galloway JM, Goley A. Scientific statement: socioecological determinants of prediabetes and type 2 diabetes. *Diabetes Care.* 2013; 36: 2430-2439.
12. Hill-Briggs F, Adler NE, Berkowitz SA, Chin MH, Gary-Webb TL. Social Determinants of Health and Diabetes: A Scientific Review. *Diabetes Care.* 2021; 44: 258-279.
13. Golden SH, Brown A, Cauley JA. Health disparities in endocrine disorders: biological, clinical, and nonclinical factors. *Endocrine Society scientific statement. J Clin Endocrinol Metab.* 2012; 97: 1579-1639.
14. U.S. Department of Health and Human Services. Section IV: Advisory Committee findings and recommendations. In *The Secretary's Advisory Committee on National Health Promotion and Disease Prevention Objectives for 2020.*
15. U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. *Healthy people 2020: social determinants of health.* Accessed 25 October 2020.
16. World Health Organization. *Diabetes 2021.*
17. Center for Disease Control and Prevention. *National Diabetes Statistics Report 2020: Estimates of Diabetes and Its Burden in the United States 2020.*
18. Choudhury S, Das SK, Hazra A. Survey of knowledge-attitude-practice concerning insulin use in adult diabetic patients in eastern India. 2014; 46: 425-430.
19. Nurumal MS, Jamaludin TSS, Mohammad NM, Hasan MKC, Win KK. A review on knowledge of diabetes and practice of medication adherence among people living with diabetes mellitus. *Int J Care Scholars.* 2020; 3: 45-54.